



Prevalence of Bovine *Trypanosomosis* and Associated Risk Factors in Jima Geneti District of the Horo Guduru Wollega Zone in Ethiopia

Dinaol Tolawak¹, Kassahun Berrie², Mahendra Pal^{3*}

¹Department of Veterinary Science, Ambo University, Ambo, Ethiopia

²Department of Veterinary Pharmacy, University of Gondar, Gondar, Ethiopia

³Narayan Consultancy on Veterinary Public Health and Microbiology- Anand-388001, Gujarat, India

Abstract

Introduction: Parasitic diseases are significant causes of morbidity and mortality in humans and animals throughout the world.

Methods: A cross-sectional study was performed in Ethiopia's Jima Geneti area of the Horo Guduru Wollega Zone from December 2017 to April 2018. The prime goal of the study was to determine the prevalence of trypanosomosis in cattle and assess potential risk factors. Buffy coat and thin blood smears were employed to identify the trypanosome species. The research district and kebeles were chosen using a purposive selection strategy, while cattle were selected by a simple random sample method.

Results: In total, 25 of the 384 investigated cattle were infected with trypanosomes, leading to an overall prevalence of 6.5%. Biqiltu Qidame town had a high prevalence of 10.4%, while Adileqa Tuluchali had a low prevalence of 3.5%. The study area had the highest prevalence of *Trypanosoma congolense* (60%), followed by *Trypanosoma vivax* (28%) and *Trypanosoma brucei* (12%). The prevalence of trypanosomosis in cattle was statistically significant ($P < 0.05$) with age, body condition, and packed cell volume (PCV) in the study area.

Conclusion: Bovine trypanosomosis was prevalent in the researched area that had a severe impact on livestock production. To increase the livestock's health and production in the study area, strategic disease prevention and control programs are necessary.

Keywords: Buffy coat, Cattle, Ethiopia, Thin blood smear, Trypanosomosis, Wollega

Received: December 5, 2021, Accepted: December 12, 2021, ePublished: January 1, 2022

Introduction

Agriculture is the world's most populous industry, employing 42% of the world's population and more than 50% of the population of developing countries (1). Livestock is the sub-sector of agriculture. Agriculture is the mainstay in Ethiopia similar to other developing countries. There are 55.03 million cattle, 27.35 million sheep, 28.16 million goats, 1.96 million horses, 6.95 million donkeys, 0.36 million mules, 1.1 million camels, and 51.35 million poultry in Ethiopia's livestock population (2).

Protozoan diseases such as toxoplasmosis, leishmaniasis, giardiasis, trypanosomosis, babesiosis, amoebiasis, and sarcocystosis are reported from many countries of the world (3,4). Among these, trypanosomosis is a serious disease in domestic livestock that causes a significant negative impact on food production and economic growth in many parts of the world (5), particularly in sub-Saharan Africa (6). There is a widespread trypanosomosis among the domestic livestock in Ethiopia's western, southern, and south-western lowlands, along with the

connected river systems (7).

The most important species affecting cattle in Ethiopia are *Trypanosoma congolense*, *Trypanosoma vivax*, and *Trypanosoma brucei* (8). Trypanosomosis is biologically transmitted by the tsetse fly, namely, *Glossina* species (3,9). The susceptibility of cattle to trypanosomosis is based on breed, age, habitat, previous exposure, and health status (10). In the pathogenesis, a local inflammatory response is called a chancre (11). There are four major stages in the life cycle of African trypanosomes. The procyclic form, epimastigote form, and metacyclic form all develop in tsetse, while the bloodstream form is found in the mammalian host (12).

The wet blood smear, thick and thin smears, buffy coats, and packed cell volume (PCV) techniques are used for the diagnosis of trypanosomosis (13,14). Control of parasites with chemotherapeutic and chemoprophylactic agents has the double effect of limiting losses caused by the infection and eliminating the transmissible trypanosome reservoir (15). To improve the welfare and security of rural communities, particularly in Ethiopia,



rapid methods for assessing risk and diagnosing urgent problems are needed for the control of animal diseases. Thus, the objective of this study is to determine the prevalence of bovine trypanosomosis and its risk factors in the Jima Geneti district of Horo Guduru Wollega Zone.

Materials and Methods

Study Area

The study was conducted in the Jima Geneti district of the Horo Guduru Wollega Zone of Oromia Regional State, Ethiopia from December 2017 to April 2018. The total livestock population in the area was 351 295 heads of cattle. A mixed crop and livestock farming system is the mode of agriculture in the district (16). The district is located about 250 km West of Addis Ababa, with geographical coordinates of 09°29'N and 37°26'E. Temperatures range from 14.9°C to 27°C monthly. The altitude of the area ranges from 1700 to 2350 above mean sea level. The area has one long rainy season extending from March to mid-October with annual rainfall ranging from 1000 to 2400 mm (17).

Study Animals and Study Population

The animals sampled for this study were local zebu cattle (*Bos indicus*) which were kept under an extensive management system (Figure 1). The cattle of both genders and all ages found in the study district were considered as study populations. All animals were grouped into three age categories, including those less than 3 years since 6 months as young, 3-5 years as an adult, and above 5 years old. The age was estimated by using dentition patterns, including teeth eruption and wear (18). In addition, the body condition was classified into three score categories of two, five, and eight representing poor, medium, and good, respectively (19).

Study Design and Sampling Technique

A cross-sectional study was conducted in December 2017 and April 2018. A purposive sampling technique was applied to select the study district and kebeles (Adileqa, Tuluchali, Biqiltu Qidame town, Damu Gembo, Gudetu Geneti, and Hareto town) based on the geographical location and their accessibility, while a simple random sampling method was used to select individual animals.

Sample Size Determination

The sample size was calculated by using a 95% confidence interval, 5% desired absolute precision, and since there was no previous study conducted in the area, 50% expected prevalence of trypanosomosis via the following formula (20).

$$N = \frac{1.96^2 P \exp(1 - P \exp)}{d^2}$$

where N , p , and d represent the required sample size, expected prevalence, and desired precision, respectively. Accordingly, the calculated sample size was 384. The number of sampled animals was not proportionately distributed but based on the abundance of study animals for each kebele.

Blood Sample Collection and Laboratory Procedure

The parasitological diagnostic tests were used based on the study purpose (21). In brief, the blood was collected from an ear vein into heparinized capillary tubes and transferred onto glass slides to make the blood smears. Each capillary tube was filled to its last third, sealed with the crystal sealant one end, and immediately centrifuged in a microhematocrit centrifuge for 5 minutes at 10 000 rpm. The PCV was determined after centrifugation, and the




Figure 1. Study Area Map Designed Using ArcMap 10.3

anemic animals were considered to have PCV < 24%. The uppermost layers of red blood cells from each specimen were removed, and the Buffy coat was placed onto a microscope slide and examined under a phase-contrast microscope using a ×40 objective lens for the presence of trypanosomes. The thick and thin blood smears were stained with Giemsa and examined under a light microscope using a ×100 oil immersion objective lens.

Data Analysis

The data were stored in a Microsoft 2013® excel spreadsheet and analyzed using STATA (22). Pearson’s Chi-square was used to evaluate the association between the prevalence of the disease with related risk factors. The confidence level was held at 95%, and the significance level was at $P < 0.05$.

Results

In the current study, the overall prevalence of trypanosomiasis was recorded at 6.5%, and the highest prevalence of 10.4% was observed in the Biqiltu Qidame town (Table 1).

Trypanosoma congolense, *Trypanosoma vivax*, and *Trypanosoma brucei* were the species of *Trypanosoma* identified by Giemsa-stained thin blood smear examination among which *T. congolense* was found to be a highly prevalent *Trypanosoma* species in the study area (Figure 2).

The anemic animals, old age, and poor body condition were statistically significant ($P < 0.05$) at a higher risk of acquiring trypanosomiasis (Table 2).

Discussion

The overall prevalence of trypanosomiasis in cattle kept under an extensive management system was 6.5% in the current study, which is in agreement with the prevalence of 5.5% in Didessa woreda, Oromiya region (23), 6.9% in Lalo Kile district of Kelem Wollega Zone (24), and 6.3% in Bako Tibe district of West Showa and Gobu Seyo districts of West Wollega Zone (25).

Moreover, almost similar prevalence of 8.5% (26) and 13.4% (27) were reported from the Sasiga and Diga districts of East Wellega and Gawo Dale districts, respectively. Compared to the present observations, the lower prevalence of 0.9%, 1.2%, and 2.1% were reported from Addisamba, Amarit, district of West Gojjam administrative zone, and Amhara region, respectively (28,29). Conversely, higher prevalence of trypanosomiasis 25.8% and 40% were reported in the Asosa district and Wolyta and Dawero zones, respectively (30,31). Those differences in the prevalence rate reported by various researchers might be due to variations in the agro-ecology, sampling season, vector infection rate, animal susceptibility, and practice of trypanocidal drug use and fly control operations which may have an impact on the

Table 1. The Overall Prevalence of Bovine Trypanosomiasis in the Study Areas

Kebeles	No. of Examined Animals	No. of Positive Animals	Prevalence (%)
Adileqa Tuluchali	57	2	3.5
Hareto town	111	4	3.6
Damu Gembo	42	2	4.8
Gudetu Geneti	78	7	9
Biqiltu Qidame town	96	10	10.4
Total	384	25	6.5

Table 2. Prevalence of Bovine Trypanosomiasis as per Different Risk Factors

Variables	Categories	No. of Examined Animals	No. of Positive Animals	Prevalence (%)	χ^2	P Value
Age	Young	116	2	1.7	6.76	0.03
	Adult	111	7	6.3		
	Old	157	16	10.2		
Body condition	Good	64	0	0	6.27	0.04
	Medium	129	8	6.2		
	Poor	191	17	8.9		
PCV	Non-anemic	176	0	0	22.63	0.00
	Anemic	208	25	12		
Gender	Male	166	9	5.4	0.57	0.53
	Female	218	16	7.3		

Note. PCV: Packed cell volume.




Figure 2. Prevalence of *Trypanosoma* Species

epidemiological situations of the disease (32,33).

In the present study, a higher prevalence of trypanosomiasis 10.4% in Biqiltu Qidame town was detected compared to 3.5% in Adileqa Tuluchali kebele. This finding is in line with 10.2% and 2.5% prevalence of trypanosomiasis in Haro Gudisa and Bila town of Gudeya Bila district, western Ethiopia, respectively. This may be due to the similarity of the addresses in the climate, altitude, and vegetation (34).

Based on the findings of this study, the majority of infections were caused by *T. congolense* (15/25, 60%),

followed by *T. vivax* (7/25, 28%) and *T. brucei* (3/25, 12%). The higher infection rate with *T. congolense* in the study area is in agreement with 58.5% *T. congolense* and 32.2% *T. vivax* reported in the southwest of Ethiopia (7). Such a high prevalence of *T. congolense* may be due to the presence of a biological vector (*Glossina*), and it is mainly confirmed in the blood, while *T. vivax* and *T. brucei* could also invade the tissues (35).

In the current study, the prevalence of trypanosomosis in old (10.2%) and adult (6.3%) animals were higher than in young (1.7%) ones based on the chi-square analysis, and there was a statistically significant seroprevalence of trypanosomiasis with age ($P=0.03$). However, this finding contradicts the prevalence of 13.8% in young, 9.4% in adults, and 4% in old animals in and around Nekemte Areas, East Wollega Zone, Ethiopia (36). In this case, older animals travel long distances for feed and draught power, as well as for harvesting crops, and this could expose them to high tsetse fly challenges (37). There is also evidence that *T. congolense* infection is a chronic disease that increases with aging (38).

In the present study, the prevalence of trypanosomosis in poor body condition (8.9%) was higher than in good body condition (0%) and there was statistically significant seroprevalence of trypanosomosis with body condition ($P=0.04$). Cattle with good body conditions were aparasitaemic for bovine trypanosomosis in this study. The absence of *Trypanosoma* infection in the good body condition of animals was because well-nourished animals have a good level of immunity and are in a better position to resist infection. Moreover, there is a highly rare possibility of the re-establishment of infection in animals with good body conditions (25). This is almost similar to the prevalence of 10.7%, 2.3%, and 1.6% in poor, medium, and good body conditions, respectively, in Mao Komo special woreda, Benishalgul Gumuz (39). This might be attributed to the reduced resistance of those animals having poor body conditions or related to the progressive weight loss arising from the debilitating nature of the disease itself (40).

In this study, the prevalence of trypanosomosis was higher in anemic animals (12%) compared to non anemic animals (0%). based on the PCV results. Animals with PCV less than 24% were considered to be anemic (41). Because the development of anemia is the most reliable indicator of *Trypanosoma* infection, even though it also interferes with concurrent diseases and nutritional factors (42,43).

Conclusion

The findings of the present study indicated that bovine trypanosomosis is a prevalent parasitic disease with an overall prevalence of 6.5%, and *T. congolense*, *T. vivax*, and *T. brucei* species were identified in the Jima Geneti district. Higher prevalence and statistical significance were observed in anemic, old age, and poor body conditions

with trypanosomosis. Biqiltu Qidame town had the highest prevalence (10.4%) among the studied kebeles. The disease could be a constraint to livestock production in the study area. Therefore, regular screening of the disease followed by early treatment of positive animals with trypanocidal drugs is necessary. An integrated tsetse control strategy should be implemented in the area. Finally, awareness creation about the economic importance of the disease and its vectors for the stakeholders is crucial.

Acknowledgments

The authors would like to acknowledge the University of Gondar for facility support and Wollega University for the provision of laboratory facilities. We dedicate our research paper to all the scientists who did pioneer work in the field of trypanosomosis.

Author Contributions

Dinaol Tolawak: Conceptualization, methodology, investigation, data curation, formal analysis, and writing-original draft.

Kassahun Berrie: Conceptualization, methodology, supervision, writing-review, and editing.

Yagoob Garedaghi: Writing-reviewing, and editing.

Mahendra Pal: Writing, reviewing, and editing.

Data Availability Statement

The data collected and used to support this article can be offered by the first author upon request.

Conflict of Interests

The authors declare that they have no conflict of interests.

Ethical Issues

Nil.

Funding

This manuscript received no financial support.

References

1. Upton M. The Role of Livestock in Economic Development and Poverty Reduction. Food and Agriculture Organization of the United Nations (FAO); 2004
2. CSA. Federal Democratic Republic of Ethiopia, Central Statistical Agency. Agricultural sampling Survey: Livestock and Livestock characteristics. 505 statistical Bulletin, Addis Ababa, Ethiopia; 2013.
3. Pal M. Zoonoses. 2nd ed. Jaipur, India: Satyam Publishers; 2007.
4. Pal M, Gutama KP, Steinmetz CH, Dave P. Leishmaniasis: an emerging and re-emerging disease of global public health concern. *Am J Infect Dis*. 2022;10(1):22-5. doi: [10.12691/ajidm-10-1-4](https://doi.org/10.12691/ajidm-10-1-4).
5. Taylor MA, Coop RL, Wall R. *Veterinary Parasitology*. Oxford: Blackwell Pub; 2007.
6. Cecchi G, Mattioli RC, Slingenbergh J, de la Rocque S. Land cover and tsetse fly distributions in sub-Saharan Africa. *Med Vet Entomol*. 2008;22(4):364-73. doi: [10.1111/j.1365-2915.2008.00747.x](https://doi.org/10.1111/j.1365-2915.2008.00747.x).
7. Abebe G. Trypanosomosis in Ethiopia. *Ethiop J Biol Sci*. 2005;4(1):75-121. doi: [10.4314/ejbs.v4i1.39017](https://doi.org/10.4314/ejbs.v4i1.39017).
8. Ayisheshim A, Abegaze S, Derso S, Shewatalek M, Hailemariam H, Shiret B, et al. Review on bovine trypanosomosis in Ethiopia. *Eur J Biol Sci*. 2015;8(1):1-7.
9. Girma K, Meseret T, Tilahun Z, Haimanot D, Firew L, Tadele K, et al. Prevalence of bovine trypanosomosis, its vector density and distribution in and around Arbaminch, Gamo Gofa Zone, Ethiopia. *Acta Parasitol Glob*. 2014;5(3):169-76.

- doi: [10.5829/idosi.apg.2014.5.3.859](https://doi.org/10.5829/idosi.apg.2014.5.3.859).
10. Orenge CO, Munga L, Kimwele CN, Kemp S, Korol A, Gibson JP, et al. Trypanotolerance in N'Dama x Boran crosses under natural trypanosome challenge: effect of test-year environment, gender, and breed composition. *BMC Genet.* 2012;13:87. doi: [10.1186/1471-2156-13-87](https://doi.org/10.1186/1471-2156-13-87).
 11. World Health Organization (WHO). Control and Surveillance of Human African Trypanosomiasis: Report of a WHO Expert Committee. WHO; 2013.
 12. WHO. Trypanosomiasis, Human African (Sleeping Sickness). 2012. Available from: <http://www.who.int/mediacentre/factsheets/fs259/en/>.
 13. World Organization for Animal Health (OIE). Trypanosomosis: Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. France: OIE; 2013.
 14. Yusuf AB, Abubakar A, Musa UB, Haruna MK, Garba HA, Maigari AK, et al. Surveillance for tsetse and trypanosomosis in Bagudo local government area north-western Nigeria. *IOSR J Agric Vet Sci.* 2015;8(7):43-8. doi: [10.9790/2380-08734348](https://doi.org/10.9790/2380-08734348).
 15. Welburn SC, Beange I, Ducrotoy MJ, Okello AL. The neglected zoonoses--the case for integrated control and advocacy. *Clin Microbiol Infect.* 2015;21(5):433-43. doi: [10.1016/j.cmi.2015.04.011](https://doi.org/10.1016/j.cmi.2015.04.011).
 16. JGWAO. Annual progress report. Jimma Geneti worda Agricultural Office. 2017.
 17. Olana BT. People and Dams: Environmental and Socio-Economic Changes Induced by a Reservoir in Fincha'a Watershed, Western Ethiopia. Netherland: Wageningen University; 2006.
 18. de Lahunta A, Habel RE. Applied Veterinary Anatomy. Philadelphia: WB Saunders Company; 1986. p. 330.
 19. Nicholson MJ, Butterworth MH. A Guide to Condition Scoring of Zebu Cattle. Addis Ababa: ILCA; 1986. p. 235.
 20. Thrusfield M. Veterinary Epidemiology. 3rd ed. Blackwell Science Ltd; 2008. p. 233-50.
 21. Lou J, Yu Y, Dia F. Laboratory test for diagnosis of parasitic disease; *Radiol.Parasit. Dis.* 2017; 6-46. doi:[10.1007/1978-94-024-0911-6-6](https://doi.org/10.1007/1978-94-024-0911-6-6).
 22. Stata Version 11. Texas, USA: Stata Corporation; 2009.
 23. Bekele N, Kebede A, Mulatu E. Prevalence of bovine trypanosomosis in Didessa woreda, Oromia region, Ethiopia. *J Vet Sci Technol.* 2018;9(1):503. doi: [10.4172/2157-7579.1000503](https://doi.org/10.4172/2157-7579.1000503).
 24. Efrem D, Bashatu F, Bacha B, Addisalem H, Misgana D. Prevalence of bovine trypanosomosis in Lalo Kile district, Kelem Wollega Zone, Oromia regional state, western Ethiopia. *Acta Parasitol Glob.* 2013;4:34-40.
 25. Abera Z, Fekadu M, Kabeta T, Kebede G, Mersha T. Prevalence of bovine trypanosomosis in Bako Tibe district of West Shoa and Gobu Seyo districts of West Wollega Zone, Ethiopia. *Eur J Biol Sci.* 2014;6(3):71-80.
 26. Tafese W, Melaku A, Fentahun T. Prevalence of bovine trypanosomosis and its vectors in two districts of East Wollega Zone, Ethiopia. *Onderstepoort J Vet Res.* 2012;79(1):E1-4. doi: [10.4102/ojvr.v79i1.385](https://doi.org/10.4102/ojvr.v79i1.385).
 27. Waktole TE. Studies on Bovine Trypanosomosis and Therapeutic Efficacy of Selected Trypanocidal Drugs in Birbir Valley of Gawo-Dalle District, West Oromia. Ethiopia: Addis Ababa University, Faculty of Veterinary Medicine Debre Zeit; 2008. p. 26-30.
 28. Addisalem HB, Tafere CA, Beshatu FW, Asamnew TM. Prevalence of bovine trypanosomosis in Addisamba and Amarit district of West Gojjam Zone, Amhara regional state. *Am J Sci Res.* 2012;7(3):112-7.
 29. Ayana M, Tesfaheywet Z, Getnet F. A cross-sectional study on the prevalence of bovine trypanosomosis in Amhara region, Northwest Ethiopia. *Livest Res Rural Dev.* 2012;24(8):1-5.
 30. Miruk A, Hagos A, Yacob HT, Asnake F, Basu AK. Prevalence of bovine trypanosomosis and trypanocidal drug sensitivity studies on *Trypanosoma congolense* in Wolyta and Dawero zones of southern Ethiopia. *Vet Parasitol.* 2008;152(1-2):141-7. doi: [10.1016/j.vetpar.2007.12.007](https://doi.org/10.1016/j.vetpar.2007.12.007).
 31. Tesfaye D, Ibrahim N. Prevalence of bovine trypanosomosis in Assosa district of Benishangul Gumuz regional state, Ethiopia. *Adv Biol Res (Rennes).* 2017;11(1):13-7.
 32. Geiger A, Ponton F, Simo G. Adult blood-feeding tsetse flies, trypanosomes, microbiota and the fluctuating environment in sub-Saharan Africa. *ISME J.* 2015;9(7):1496-507. doi: [10.1038/ismej.2014.236](https://doi.org/10.1038/ismej.2014.236).
 33. Majekodunmi AO, Fajinmi A, Dongkum C, Picozzi K, Thrusfield MV, Welburn SC. A longitudinal survey of African animal trypanosomiasis in domestic cattle on the Jos Plateau, Nigeria: prevalence, distribution and risk factors. *Parasit Vectors.* 2013;6(1):239. doi: [10.1186/1756-3305-6-239](https://doi.org/10.1186/1756-3305-6-239).
 34. Abdeta T, Kebede A, Dugassa J, Abera Z. Prevalence of bovine trypanosomosis in Gudeya Bila district, East Wollega Zone of Oromia regional state, western Ethiopia. *Afr J Basic Appl Sci.* 2016;8(6):336-42.
 35. Fayisa G, Mandefro A, Hailu B, Chala G, Alemayehu G. Epidemiological status and vector identification of bovine trypanosomosis in Didesa district of Oromia regional state, Ethiopia. *Int J Nutr Food Sci.* 2015;4(3):373-80.
 36. Gameda F. Prevalence of bovine trypanosomosis in and around Nekemte areas, East Wollega Zone, Ethiopia. *Open Access Libr J.* 2015;2(5):68408. doi: [10.4236/oalib.1101521](https://doi.org/10.4236/oalib.1101521).
 37. Batu G, Abera Z, Tadesse NN, Wakgari M, Moti A. Prevalence of bovine trypanosomosis in Gimbi district, West Wollega, Western Oromia of Ethiopia. *Open Access J Vet Sci.* 2017;3(5):1-9.
 38. McDermott J, Woitag T, Sidibé I, Bauer B, Diarra B, Ouédraogo D, et al. Field studies of drug-resistant cattle trypanosomes in Kéné Dougou province, Burkina Faso. *Acta Trop.* 2003;86(1):93-103. doi: [10.1016/s0001-706x\(03\)00019-6](https://doi.org/10.1016/s0001-706x(03)00019-6).
 39. Haile G, Gizaw O. Cross sectional study on prevalence of bovine trypanosomosis and associated risk factors in Mao Komo special woreda, Benishangul Gumuz, Western Ethiopia. *J Parasitol Vector Biol.* 2018;10(4):45-50.
 40. Radostits OM, Gay CC, Hinchcliff KW, Constable PD. *Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats.* 10th ed. Saunders Ltd; 2007. p. 1531-40.
 41. World Organization for Animal Health (OIE). Trypanosomosis (Tsetse- Transmitted): Terrestrial Manual. Paris, France: OIE; 2008. p. 654.
 42. Garedaghi G, Firouzvand Y, Heikal Abadi M. Assessment of *Neospora caninum* seroprevalence in buffalo in Tabriz city, north-west of Iran. *Buffalo Bull.* 2017;36(2):379-84.
 43. Abebe G. Trypanosomosis in Ethiopia. Debre Zeit: Faculty of Veterinary Medicine, Addis Ababa University; 2005. p. 18-20.