



Diversity of Intestinal Parasites in Ostriches *Struthio camelus* Linnaeus, 1758 by Using Routine Parasitological Techniques and PCR in Iraq

Zainab A. Makawi¹ , Maryam M Al-Khaiat¹ , Hiba Mohammed Jihad¹

¹Iraq Natural History Research Center and Museum, University of Baghdad, Baghdad, Iraq

Abstract

Introduction: Ostrich farming has emerged as a new livestock industry in Iraq, but scientists lack sufficient information on health concerns, including intestinal parasites that cause significant production losses and financial instability over extended periods.

Methods: Researchers collected 150 fecal samples from ostriches that dwelled in central and southern Iraq for microscopic examination of intestinal parasite occurrence.

Results: The six parasite species included *Entamoeba* sp., which made up 26.66% of the population, and *Cryptosporidium* sp. at 11.33%, *Ascaridia galli* at 10%, *Giardia* sp. at 4.6%, *Raillietina* sp. at 2%, and *Trichostrongyl*. Molecular analysis was performed on a subset of positive samples because *Entamoeba* sp. is highly prevalent. PCR amplification of the 18S rRNA gene revealed fragments of approximately 579 bp for *Entamoeba struthionis* (IDs: PV019353.1, PV019354.1), *Entamoeba polecki* (IDs: PV019355.1, PV019356.1), and *Entamoeba* sp. (ID: PV019357.1), the first time in Iraq. The NCBI database now has these sequences.

Conclusion: The current study concluded that molecular diagnostics in ostrich health management are crucial for early detection, precise treatment, and improved productivity. Regular monitoring is recommended to promote sustainable ostrich farming in Iraq.

Keywords: Intestinal parasite, Iraq, PCR reaction, Ostriches, 18S rRNA

Received: December 12, 2025, Revised: December 20, 2025, Accepted: December 31, 2025, ePublished: March 8, 2026

Introduction

Ostriches, one of the world's strongest and most resilient birds, can be raised outside of Africa in both northern and southern climates. Climate-related issues, however, may affect the profitability of ostrich farming. Feathers, leather, and meat are just a few of the goods produced by the ostrich industry (1). Organisms that reside within or on another species are known as parasites (2). In addition to being essential for maintaining ecological balance, wild animals are valuable human companions. In addition to their crucial role in sustaining ecological balance, wild animals serve as significant companions to humans.

Over the past six decades, considerable ecological changes and habitat degradation have influenced the ostrich industry (3). Birds serve as hosts for numerous parasite species, which have adapted to live within them. The consumption of parasite eggs or oocysts does not always result in infection, and parasitism may not always present as a clinical disease (4). Birds serve as essential hosts for generalist parasites because they inhabit various environments and function as reservoirs for these parasites (5). The proper use of anthelmintics requires veterinary guidance because incorrect application can lead to drug resistance and persistent parasite infections (6). The ostrich is particularly susceptible to protozoan parasites,

including *Entamoeba struthionis* (7). The prevalence of animal parasites depends on multiple factors, which include host immune defenses, environmental conditions, habitat characteristics, and species traits (8,9). Wildlife species require monitoring systems for parasitic infections to prevent disease transmission among species and protect their survival. The presence of *Strongyloides* and *Toxocara* parasites in animal habitats near human settlements increases the risk of zoonotic disease (10–12).

The parasites that cause cestode infections, which lead to tissue necrosis and vascular obstruction, pose a risk to animals living near populated areas. Parasites harm host tissues by physically entering host tissues when their larvae penetrate during the reproductive or establishment phases (13). Wild bird species are essential components of ecosystem and biodiversity protection. Wild birds perform essential functions as natural pest controllers, seed dispersers, and pollinators (14, 15). The reduction of biodiversity accelerates disease spread, creating more opportunities for zoonotic diseases to affect both domestic animals and people while simultaneously leading to the emergence of new disease outbreaks. This is especially relevant for species that regulate disease and pest populations. The established factors create a destructive feedback loop that threatens worldwide



health protection systems (16, 17). Wildlife conservation initiatives should establish new populations and protect existing populations through habitat conservation to safeguard the genetic diversity of animal populations and endangered species. The practice of captivity helps protect birds from specific dangers, but it creates new stressful conditions that increase their susceptibility to parasites and health problems (18). Regular parasitological testing, together with advanced diagnostic methods, is essential for reducing adverse outcomes associated with parasitic infections. The techniques enable doctors to detect diseases in their early stages, thereby allowing them to develop specific treatment strategies (19).

The research focused on the scientific study of intestinal parasite infections affecting ostriches. The particular objectives of the study were:

1. Through microscopic examination and molecular analysis, researchers will identify all parasite species that affect ostriches.
2. To evaluate the effectiveness of advanced diagnostic techniques for diagnosing parasitic infections in ostriches.

Materials and methods

Collection sample

The researchers collected fresh ostrich feces samples from 64 (42.66%) male and 86 (57.33%) female who lived in cages and reserves across different Iraqi locations which included Al-Mahawil (Babil), Al-Zawraa Park, Al-Adamiyah (Baghdad), and Azizia (Wasit) as shown in Figure 1 from November 2023 to July 2024, their ages

ranged from one year to over three years old. The ostrich received four samples, each weighing between 10 and 20 grams, collected every three days. The samples were stored in a sterile container in a freezer until laboratory testing. The animals were assigned identification numbers when scientists collected their samples.

Microscopic examination

At the Vertebrate Laboratory, INHM University of Baghdad, faecal samples were processed according to established coprological protocols (20-21). The process of detecting intestinal parasites involved centrifugation to concentrate samples, followed by flotation in Sheather's sugar solution and staining with 2% Lugol's Iodine to make cysts, oocysts, and eggs more visible. A calibrated ocular micrometer was used to obtain morphometric measurements, and high-resolution digital photographs were used to document the parasitic structures. Standard diagnostic methods reach their full potential when combined with PCR-based molecular tests, which identify parasite species at low concentrations and in complex fecal sample matrices.

Molecular study

PCR Amplification and DNA Extraction

A commercial DNA extraction kit (Intron Biotechnology, Korea) was used to extract genomic DNA from 50 *Entamoeba* sp.- positive fecal samples that had previously been collected and stored at -20 °C in accordance with the manufacturer's instructions. The purified DNA samples were stored at -20 °C before use.




Figure 1. Geographical distribution of ostrich farming sites sampled in central Iraq

We amplified the 18S rRNA gene for molecular characterization using the Verweij *et al.* protocol (22). With the primer pair: Forward 5'-GTTGATCCTGCCAGTATTATATG-3' and Reverse 5'-CACTATTGGAGCTGGAATTAC-3'. Ten representative samples were selected from the extracted DNA and subjected to PCR amplification. The 25 μ L total volume of the PCR reaction was composed of 12.5 μ L of GoTaq Green Master Mix (Promega, USA), 1 μ L of each forward and reverse primer (10 pmol/ μ L), 1.5 μ L of template DNA, and 9 μ L of nuclease-free water. The mixture was gently vortexed and briefly centrifuged, then subjected to PCR amplification.

The PCR products were analyzed by electrophoresis on a 1.5% agarose gel prepared with 1 \times TBE buffer. A GelDocTM XR+imaging system (Bio-Rad, USA) was used to visualize the gel under UV light after staining with ethidium bromide (0.5 μ g/ml). The size marker was a 100-bp DNA ladder (Promega, USA). According to Verweij *et al.* (22), the *Entamoeba* 18S rRNA fragment was expected to be about 585 bp in size. In this study, the PCR products migrated near 579 bp, consistent with the anticipated size. Slight deviations in length (\pm 5–10 bp) were due to variations in gel concentration and electrophoretic conditions.

Sequencing of DNA and phylogenetic tree

To ascertain the molecular identity of *Entamoeba* species, DNA sequencing was performed on 10 positive samples using PCR-amplified *Entamoeba* spp. DNA. The amplified segment of the 18S rRNA gene was purified using the Strata Prep PCR Purification Kit. The UPGMA [unweighted pair group method with arithmetic mean] method was used to infer the evolutionary history. A phylogenetic tree and the GenBank database were used to obtain accession numbers for local *Entamoeba* species in Iraq, and bioinformatics for Ostrich *Entamoeba*.

The purified 18S rRNA PCR amplicons were sent to Macrogen Inc. for Sanger sequencing. To acquire accession numbers, the resulting *Entamoeba* spp. nucleotide sequences were entered into the NCBI GenBank database. ClustalW was used to align the sequences with reference sequences obtained from GenBank for subsequent analysis. MEGA version 6 was used to infer phylogenetic relationships using conventional evolutionary models. *Species identification of Entamoeba isolates was verified by BLAST against reference sequences* in the NCBI database.

Statistical Analysis

A statistical analysis was performed to assess the relationship between geographical location, and the results showed a slight variation in the spread. However, it was not statistically significant at the 0.05 level ($\chi^2 = 6.47$, $df = 3$, $P = 0.09$).

Results

This study was conducted at multiple sites across central Iraq, including Al-Mahawil (Hilla), Al-Zawraa Park, Al-Adamiyah (Baghdad), and Azizia (Wasit), from November 2023 to July 2024. Of 150 fecal samples, 85 (56.66%) were positive for intestinal parasites; the difference was not statistically significant at the 0.05 level (Table 1). Several different intestinal parasites were identified using the feces of ostriches based on their morphological traits and size included six species of intestinal parasite: *Entamoeba* sp (26.66%), *Cryptosporidium* sp. (11.33%), *Ascaridia galli* (10%), *Giardia* sp (4.6%), *Raillietina* sp (2%) and *Trichostrongylus tenuis* (2%), (Figure 2, Tables 2 and 3).

Molecular results

Gel electrophoresis on 1.5% agarose showed clear, individual amplicon bands approximately 579 bp in length in all positive samples, consistent with the expected 585 bp fragment of the *Entamoeba* 18S rRNA gene (Figure 3).

This confirms the successful molecular identification of *Entamoeba* sp. Minor size variation (520–585 bp) across runs was attributable to electrophoresis conditions and marker calibration, rather than sequence differences.

DNA sequencing results

The current study identified 50 samples suspected of containing *Entamoeba* sp. Ten were subjected to PCR, and five were sent for phylogenetic analysis and sequencing. MEGAX software was used to analyse the sequenced samples (23). The similarity percentage between the positive isolates and the NCBI isolates was calculated. The accession numbers for the local ostrich *Entamoeba* species were obtained by uploading the sequences to NCBI GenBank.

Three species were identified based on the 18S rRNA gene, as shown in Table 4: *Entamoeba struthionis* (ID PV019353.1, ID: PV019354.1), *Entamoeba polecki* (ID: PV019355.1, ID: PV019356.1), and *Entamoeba* sp (ID: PV019357.1).

Phylogenetic evaluation

A phylogenetic tree of *Entamoeba* species, constructed from previously identified 18S ribosomal RNA sequences and illustrated with colored triangles and corresponding accession numbers, showed that the Iraqi isolates clustered closely together. Specifically, *Entamoeba struthionis* was closely related to isolates from Spain and Cameroon, while *Entamoeba polecki* clustered with isolates from China and Vietnam. In contrast, the *Entamoeba* sp. isolate displayed the highest similarity to a Chinese strain (Diagram 1). The PCR amplification results, confirmed by electrophoresis, revealed the expected band size (Figure 3).

Discussion

This study gathered 150 ostrich droppings. Ostriches

Table 1. Total infection with intestinal parasites in Ostriches according to the location study

Location	Total Samples (n)	Positive Samples (n)	Negative Samples (n)	Prevalence (%)
Al-Mahawil (Babil)	35	17	18	50.0
Al-Zawraa Park (Baghdad)	45	32	13	70.0
Al-Adhamiyah (Baghdad)	40	24	16	60.0
Al-Azizia (Wasit)	30	12	18	40.0
Total	150	85	65	56.66


Not statistically significant at the 0.05 level ($\chi^2=6.47$, $df=3$, $P=0.09$).

Table 2. Prevalence of intestinal parasites according to family parasite in this research

Parasite	Family parasite	Stage	Parasite Group
<i>Entamoeba</i> sp	Entamoebidae	Cyst	Protozoa
<i>Cryptosporidium</i> sp	Cryptosporidiidae	Oocyst	Protozoa
<i>Ascaridia galli</i>	Ascarididae	Egg	Nematode
<i>Giardia</i> sp	Hexamitidae	Cyst	Protozoa
<i>Raillietina</i> sp	Davaineidae	Egg	Cestode
<i>Trichostrongylus tenuis</i>	Trichostrongylidae	Egg	Nematode

Table 3. Frequency distribution of intestinal parasite species among 150 clinical samples

Parasite	No. of positive	Percentage (%)
<i>Entamoeba</i> sp	40	26.66
<i>Cryptosporidium</i> sp	17	11.33
<i>Ascaridia galli</i>	15	10
<i>Giardia</i> sp	7	4.6
<i>Raillietina</i> sp	3	2
<i>Trichostrongylus tenuis</i>	3	2

**Figure 2.** A. Cyst of *Entamoeba* sp, B. *Cryptosporidium* sp oocyst, C. *Ascaridia galli* egg, D. Cyst of *Giardia* sp, E. *Raillietina* sp egg, F. *Trichostrongylus tenuis* egg

were selected using a technique that ensured samples were drawn from multiple farms to account for variation in parasite prevalence. 85 (56.66%) of these samples contained intestinal parasites. This finding offers significant new information about the scant studies on parasites in ostriches. Most parasites in these birds are found in the digestive system. The examination of ostriches revealed no health problems despite the presence of intestinal parasites, including species that commonly harm ruminants. This asymptomatic carriage could facilitate parasite spread between farms, particularly when animals are transported between facilities and countries (24). As a result, farmed ostriches may serve as reservoirs for a variety of intestinal parasites, posing a risk to humans,

other animals, and farm laborers. Contaminated feed, water, insect vectors, and human contact are all routes by which parasites spread. Regular parasite screening, controlled animal movement, good sanitation, and staff hygiene training are examples of preventive measures (25). However, there was no subtype overlap for *Cryptosporidium* spp. between domestic animals, local villagers, and *Giardia duodenalis*, suggesting a limited role in direct zoonotic transmission. The discovery of zoonotic protozoa in domestic animals underscores the need for heightened caution when interacting with them.

Seasonal fluctuations, sample size, local methods of parasite control, and regional and climatic factors are likely to influence the prevalence of parasites in ostriches




Figure 3. PCR amplification of the *Entamoeba* 18S rRNA gene (~579 bp) is demonstrated by agarose gel electrophoresis (1.5%). Lane M is a 100 bp DNA ladder; Lanes 1 through 5 are positive samples with clear single bands; the remaining lanes are negative controls




Diagram 1. Phylogenetic tree analysis of *Entamoeba* species with a focus on the 18S rRNA gene

Table 4. Homology of Local *Entamoeba* sp. Isolates with GenBank References

No.	Local Isolate ID	Closest Species (NCBI)	GenBank Accession No.	Country	% Identity	Bootstrap Value
1	PV019353.1	<i>Entamoeba struthionis</i>	AB851494.1	Spain	99%	99.56%
2	PV019354.1	<i>Entamoeba struthionis</i>	KF646239.1	Cameroon	99%	99.12%
3	PV019355.1	<i>Entamoeba polecki</i>	PP735780.1	China	100%	100%
4	PV019356.1	<i>Entamoeba polecki</i>	PP735780.1	Vietnam	99%	100%
5	PV019357.1	<i>Entamoeba</i> sp.	MT798826.1	China	99%	99.29%

(26). In this study, two *Entamoeba* species—*Entamoeba struthionis* and *Entamoeba polecki*—previously identified in ostriches were reported for the first time in Iraq. The researchers documented the initial observation of *Entamoeba* with a single-nucleus mature cyst in an

ostrich through their study (27), which confirmed results from Ponce-Gordo *et al.* (28) who identified *Entamoeba* species in *Struthio camelus*. Analysis of full-length short ribosomal RNA gene sequences revealed substantial genetic differences between this species and other

Entamoeba species. Simonidou *et al.* (29-31) reported that *Balantidium coli*, along with *E. struthionis*, appeared for the first time in ostriches, which Greece had never seen before. Scientists used molecular amplification techniques on the ITS region and partial 16S rRNA gene sequences to study different types of *Entamoeba* strains, which tend to form clusters. The research findings indicate that ostriches experience protozoan infections more frequently than helminth infections because their environment contains silt contaminated with fecal matter.

Previous research has shown that parasite prevalence depends on the sampling site, living environment, geographical conditions, and sanitation practices (32, 33). The genus *Entamoeba* exhibits considerable adaptability, as it can establish itself in the digestive tracts of diverse animal species (34-37).

Conclusion

The study focused on identifying intestinal parasites affecting ostriches (*Struthio camelus*). The researchers identified two new *Entamoeba* species that had not been detected in Iraq. The study shows that these parasites can infect people as they spread into areas where they were not previously observed. The study indicates that ostriches function as disease vectors, transmitting infections to other species through contact, even when they appear healthy. Several *Entamoeba* species pose a threat to human health. Assessing the risk of human infection from ostrich-derived species requires particular attention, given that people interact with these animals. Research on transmission methods, host range, and disease transmission to humans in Iraq requires additional epidemiological and molecular studies to be conducted by public health officials and researchers.

Authors' Contribution

Conceptualization, Data curation, Investigation, Project administration, Software, and Supervision of the research idea for this study originated with Zainab A. Makawi, together with Maryam M. Al-Khaiat and Hiba Mohammed Jihad. Zainab A. Makawi conducted the formal analysis, methodology, validation, visualization, microscopic diagnosis, genetic analysis, and writing—original draft of the manuscript. All authors approved the final manuscript for submission after completing their individual contributions to the paper.

Competing Interests

The publication of this article receives full disclosure for potential conflicts of interest from its authors.

Ethical Approval

Not relevant.

Funding

None.

References

- Pittaway T, Van Niekerk P. Horizon-scanning the ostrich industry with bibliometric indicators. *AFJARE*. 2015; 10(1): 64–71. doi:10.22004/ag.econ.200602
- Cabodevilla X, Aebischer J N, Mougeot F, Morales M B, Arroyo B. Are population changes of endangered little bustards associated with releases of red-legged partridges for hunting? A large-scale study from central Spain. *Eur J Wildl Res*. 2020;66(2):1–10. doi:10.1007/s10344-020-1366-3
- Christe P. [Loss of biodiversity: a one-way ticket to new emerging diseases?]. *Rev Med Suisse*. 2023;19(825):842–4. doi:doi:10.53738/revmed.2023.19.825.842
- Papini R, Girivetto M, Marangi M, Mancianti F, Giangaspero A. Endoparasite infections in pet and zoo birds in Italy. *ScientificWorldJournal*. 2012;2012:253127. doi:10.1100/2012/253127
- Ishtiaq F, Renner S. Bird migration and vector-borne parasite transmission. In: Santiago-Alarcon D, Marzal A, eds. *Avian Malaria and Related Parasites in the Tropics*. Springer; 2020:513-526. doi:10.1007/978-3-030-51633-8_16
- Nurchahyo RW, Ekawasti F, Haryuningtyas D, Wardhana AH, Firdausy LW, Priyowidodo D, et al. Occurrence of gastrointestinal parasites in cattle in Indonesia. *IOP Conference Series: Earth and Environmental Science*. 2021;686(1):012063. doi:10.1088/1755-1315/686/1/012063
- Aryani I, Suyadi, Hartutik H, Kuswati K. Prevalence and diversity of gastrointestinal endoparasite in Madura cattle at Papabaru, East Java, Indonesia. *Adv Anim Vet Sci*. 2022;10(10):2171–2177. doi:10.17582/journal.aavs/2022/10.10.2171.2177
- Sharma HP, Achhami B. Gastro-intestinal parasites of sympatric red panda and livestock in protected areas of Nepal. *Vet Med Sci*. 2022;8(2):568–77. doi:10.1002/vms3.651
- Maganga GD, Makouloutou-Nzassi P, Boundenga L, Maganga Landjekpo HN, Banguéboussa F, Ndong Mebaley T, et al. Gastrointestinal Polyparasitism in Bushmeat in Zadié Department in Northeast Gabon. *Vet Sci*. 2023;10(3). doi:10.3390/vetsci10030229
- Sapkota S, Regmi S, Ghimire R, Pokhrel A, Sadaula A, Shrestha BK. Transmission of gastrointestinal parasites between deer and livestock species at buffer zone located within the eastern part of Chitwan National Park, Nepal. *Res Sq*. 2021. doi:10.21203/rs.3.rs-746759/v1
- Paudel P, Subedi JR, Pathak A. Prevalence of gastrointestinal parasites among greater one-horned rhino in Chitwan National Park, Nepal. *Ann Parasitol*. 2022;68(1):159–67. doi:10.17420/ap6801.420
- Adhikari RB, Adhikari Dhakal M, Ale PB, Regmi GR, Ghimire TR. Prevalence of Intestinal Parasites in Captive Asian Elephants (*Elephas maximus* Linnaeus, 1758) in Central Nepal. *Vet Med Sci*. 2025;11(3):e70310. doi:10.1002/vms3.70310
- Mohammad ZAA, Awad AHH. Pathogenicity of two nematodes Tetramerese sp. and Microtetramerese spiralis in proventriculus of some aquatic birds from Al-Sanaf Marshes in Thi-Qar Province, South of Iraq. *Univ Thi-Qar J Sci*. 2020;7(2):713. doi:10.32792/utq/utjsci/v7i2.713
- Pigot AL, Sheard C, Miller ET, Bregman TP, Freeman BG, Roll U, et al. Macroevolutionary convergence connects morphological form to ecological function in birds. *Nat Ecol Evol*. 2020;4(2):230–9. doi:10.1038/s41559-019-1070-4
- Katumo DM, Liang H, Ochola AC, Lv M, Wang QF, Yang CF. Pollinator diversity benefits natural and agricultural ecosystems, environmental health, and human welfare. *Plant Divers*. 2022;44(5):429–35. doi:10.1016/j.pld.2022.01.005
- Yasmeen N, Jabbar A, Shah T, Fang LX, Aslam B, Naseeb I, et al. One Health Paradigm to Confront Zoonotic Health Threats: A Pakistan Prospective. *Front Microbiol*. 2021;12:719334. doi:10.3389/fmicb.2021.719334
- Lorenzo-Rebenaque L, López-Fernández S, Marco-Jiménez F, Montoro-Dasi L, Marin C, Vega S, et al. Zoonotic

- Parasites in Playgrounds in Southern Spain: A One Health Approach. *Microorganisms*. 2023;11(3). doi:10.3390/microorganisms11030721
18. Lacerda M, de Magalhães Santos WH, Lopes MC, Fonseca CS, de Carvalho MPN, da Silva Martins NR, et al. A Survey of Diseases in Different Species of Wild, Captive, and Illegally Traded Birds in Brazil. *Animals (Basel)*. 2023;14(1). doi:10.3390/ani14010025
 19. Egan S, Barbosa AD, Feng Y, Xiao L, Ryan U. The risk of wild birds contaminating source water with zoonotic *Cryptosporidium* and *Giardia* is probably overestimated. *Sci Total Environ*. 2024;912:169032. doi:10.1016/j.scitotenv.2023.169032
 20. Marcer F, Cassini R, Parisotto N, Tessarin C, Marchiori E. A Comparative Study of Mini-FLOTAC With Traditional Coprological Techniques in the Analysis of Cetacean Fecal Samples. *Front Vet Sci*. 2022;9:908486. doi:10.3389/fvets.2022.908486
 21. Molla YD, Alemu HT. New developments in diagnosis of intestinal parasites. In: *Intestinal Parasites – New Developments in Diagnosis, Treatment, Prevention and Future Directions*. IntechOpen; 2024. <https://www.intechopen.com/chapters/100845>
 22. Verweij JJ, Laeijendecker D, Brienen EA, van Lieshout L, Polderman AM. Detection and identification of entamoeba species in stool samples by a reverse line hybridization assay. *J Clin Microbiol*. 2003;41(11):5041–5. doi:10.1128/jcm.41.11.5041-5045.2003
 23. Tamura K, Stecher G, Peterson D, Filipinski A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol*. 2013;30(12):2725–9. doi:10.1093/molbev/mst197
 24. El-Sayed NM. Occurrence of some parasites in farmed ostriches (*Struthio camelus*) in Egypt. *Egyptian Veterinary Medical Society of Parasitology Journal*. 2016;12:91-100. doi:10.21608/evmspj.2016.37345
 25. Adeola A, Akande O, Adeniji O, Onihunwa J, Olaiifa O, Deji J. Gastrointestinal parasites of birds in zoological garden of University of Ilorin. *Egyptian Journal of Animal Production*. 2022;59(3):131-5. doi:10.21608/ejap.2022.151808.1045
 26. Lozano J, Almeida C, Victório AC, Melo P, Rodrigues JP, Rinaldi L, et al. Implementation of Mini-FLOTAC in Routine Diagnosis of *Coccidia* and Helminth Infections in Domestic and Exotic Birds. *Vet Sci*. 2021;8(8). doi:10.3390/vetsci8080160
 27. Ponce Gordo F, Herrera S, Castro AT, García Durán B, Martínez Díaz RA. Parasites from farmed ostriches (*Struthio camelus*) and rheas (*Rhea americana*) in Europe. *Vet Parasitol*. 2002;107(1-2):137–60. doi:10.1016/s0304-4017(02)00104-8
 28. Hao Y, Liu A, Li H, Zhao Y, Yao L, Yang B, et al. Molecular characterization and zoonotic potential of *Cryptosporidium* spp. and *Giardia duodenalis* in humans and domestic animals in Heilongjiang Province, China. *Parasites & Vectors*. 2024;17(1):155. doi:10.1186/s13071-024-06219-3
 29. Makawi ZA. Isolation and identification of *Eimeria* Schneider, 1875 species (Apicomplexa, Eimeriidae) from goats in Wasit province, Iraq. *Bull Iraq Nat Hist Mus*. 2024;18(2):431-445. doi:10.26842/binhm.7.2024.18.2.0431
 30. Martínez-Díaz RA, Herrera S, Castro A, Ponce F. *Entamoeba* sp. (Sarcomastigophora: Endamoebidae) from ostriches (*Struthio camelus*) (Aves: Struthionidae). *Vet Parasitol*. 2000;92(3):173–9. doi:10.1016/s0304-4017(00)00314-9
 31. Symeonidou I, Diakou A, Papadopoulos E, Ponce-Gordo F. Endoparasitism of Greek ostriches: First report of *Entamoeba struthionis* and *Balantioides coli*. *Vet Parasitol Reg Stud Reports*. 2019;18:100334. doi:10.1016/j.vprsr.2019.100334
 32. Alharazi T, Bamaga OA, Al-Abd N, Alcantara JC. Intestinal parasitic infection: prevalence, knowledge, attitude, and practices among schoolchildren in an urban area of Taiz city, Yemen. *AIMS Public Health*. 2020;7(4):769–77. doi:10.3934/publichealth.2020059
 33. Dos Santos Zanetti A, Malheiros AF, de Matos TA, Dos Santos C, Battaglini PF, Moreira LM, et al. Diversity, geographical distribution, and prevalence of *Entamoeba* spp. in Brazil: a systematic review and meta-analysis. *Parasite*. 2021;28:17. doi:10.1051/parasite/2021028
 34. Yagoob, G., Khaki, A., Raza, S.H.A., Abdelgayed, S.S., Kakar, M.U. Epidemiological and pathological studies on the helminthic parasites in native chickens of Tabriz city, Iran. *Genetics and Molecular Research*. 2017; 16(4):1-8. doi:10.4238/gmr16039824
 35. Khayatnouri MH, Garedaghi Y, Arbati AR, Khalili H, The Effect of Ivermectin Pour-on Administration Against Natural *Heterakis gallinarum* Infestation and its Prevalence in Native Poultry. *American Journal of Animal and Veterinary Sciences*. 2011 ;6(1): 55-58.
 36. Garedaghi Y. Identification of Immunogenic Relevant Antigens in the Excretory-secretory (ES) Products of *Ascaridia galli* Larvae. *Adv Environ Biol*. 2011 ;5(6):1120-1126.
 37. Khayatnouri M, Garedaghi Y, Khalili H, Arbati A. The Effect of Ivermectin Pour-on Administration Against Natural *Ascaridia Gallii* Infestation and its Prevalence in Native Poultry. *Adv Environ Biol*. 2011 5(7): 2002-2005.

© 2026 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.